AMENDMENTS TO THE CLAIMS

1. (Original) A method for producing carotenoids or their precursors using genetically modified organisms of the Blakeslea genus, which method comprises the following steps:

- (i) transformation of at least one of the cells,
- (ii) homokaryotic conversion of the cells obtained in step (i) to produce cells in which one or more genetic characteristics of the nuclei are all modified in an identical manner and said genetic modification manifests itself in the cells, and
- (iii) selection and reproduction of the genetically modified cell or cells,
- (iv) cultivation of the genetically modified cells,
- (v) preparation of the carotenoid produced by the genetically modified cells or the carotenoid precursor produced by said genetically modified cells.
- 2. (Original) The method according to claim 1, wherein the cells are from fungi of the Blakeslea trispora species.
- 3. (Currently amended) The method according to claim 1 or 2, wherein a vector or free nucleic acids are used in the transformation of step (i).
- 4. (Currently amended) The method according to claim 3, wherein the vector employed in the transformation (i) is integrated into the genome of at least one of the cells.
- 5. (Currently amended) The method according to claim 4, wherein the vector employed in the transformation (i) comprises a promoter and/or a terminator.
- 6. (Currently amended) The method according to any of the preceding claims 3 to 5 claim 3, wherein a vector comprising the a gpd, pcarB, pcarRA and/or ptef1 promoter and/or the a trpC terminator is employed in the transformation-(i).
- 7. (Currently amended) The method according to any of the preceding claims 3 to 6 claim 3, wherein a vector comprising a resistance gene is employed in the transformation (i).

8. (Currently amended) The method according to claim 7, wherein the vector employed in the transformation (i) comprises a hygromycin resistance gene (hph), in particular from *E. coli*.

- 9. (Currently amended) The method according to any of the preceding claims 5—8 claim 6, wherein the gpd promoter has comprises the sequence SEQ ID NO: 1.
- 10. (Currently amended) The method according to any of the preceding claims 5 8 claim 6, wherein the trpC terminator has comprises the sequence SEQ ID NO: 2.
- 11. (Currently amended) The method according to any of the preceding claims 5 8 claim 6, wherein the tef1 ptef1 promoter has comprises the sequence SEQ ID NO: 35.
- 12. (Currently amended) The method according to any of claims 6 to 11 claim 6, wherein the gpd promoter and the trpC terminator are derived from Aspergillus nidulans.
- 13. (Currently amended) The method according to any of claims 3 to 12 claim 3, wherein the vector comprises the sequence SEQ ID NO: 3.
- 14. (Currently amended) The method according to any of the preceding claims claim 1, wherein the transformation (i) is carried out using agrobacteria, conjugation, chemicals, electroporation, bombardment with DNA-loaded particles, protoplasts or microinjection.
- 15. (Currently amended) The method according to any of the preceding claims claim 1, wherein a mutagenic agent is employed in the homokaryotic conversion of step (ii).
- 16. (Original) The method according to claim 15, wherein the mutagenic agent employed is N-methyl-N'-nitronitrosoguanidine (MNNG), UV radiation or X rays.
- 17. (Currently amended) The method according to any of the preceding claims claim 1, wherein the selection is carried out by labeling and/or selecting the mononuclear cells.
- 18. (Currently amended) The method according to any of the preceding claims 1—17 claim

 1, wherein 5-carbon-5-deazariboflavin (darf) and hygromycin (hyg) or 5-fluororotate

 (FOA) and uracil and hygromycin are employed in the selection.

19. (Currently amended) The method according to any of claims 3 to 18 claim 3, wherein the vector employed in the transformation (i) includes genetic information for producing carotenoids or their precursors.

- 20. (Currently amended) The method according to any of claims 3 to 19 claim 3, wherein the vector employed in the transformation (i) includes genetic information for producing carotenes or xanthophylls.
- 21. (Currently amended) The method according to any of claims 3 to 20 claim 3, wherein the vector employed in the transformation (i) includes genetic information for producing astaxanthin, zeaxanthin, echinenone, β-cryptoxanthin, andonixanthin, adonirubin, canthaxanthin, 3-hydroxyechinenone, 3'-hydroxyechinenone, lycopene, β-carotene, α-carotene, lutein, phytofluene, bixin or phytoene.
- 22. (Currently amended) A method for providing at least one highly pure carotenoid and a foodstuff comprising carotenoid-producing organisms and at least the one carotenoid, which method comprises, after cultivation of carotenoid-producing genetically modified organisms of the Blakeslea genus according to any of claims 1 to 21 claim 1, the following steps:
 - I) removal of the biomass,
 - IA) optional washing of the biomass with a solvent in which carotenoids are not soluble, in particular water,
 - IB) sterilization and cell disruption of the biomass,
 - IC) optional drying and/or homogeneous distribution, and
 - partial extraction of the carotenoids from the disrupted biomass by means of a carotenoid-dissolving solvent and separation of said solvent from said biomass,

IIA)

1) removal of residual solvent from the carotenoid-containing biomass,

2) optional homogeneous suspension of the biomass, with a biomass solid content of > 2% and < 50%,

3) drying of the biomass or suspension for producing the foodstuff,

IIB)

- 1) crystallization of the carotenoids from the solvent used and isolation of the carotenoid crystals, in particular by filtration.
- 23. (Original) The method according to claim 22, wherein the at least one carotenoid is selected from the group consisting of carotenes and xanthophylls.
- 24. (Currently amended) The method according to claim 22 or 23, wherein the at least one carotenoid is selected from the group consisting of astaxanthin, zeaxanthin, echinenone, β-cryptoxanthin, andonixanthin, adonirubin, canthaxanthin, 3-hydroxyechinenone, 3'-hydroxy-echinenone, lycopene, β-carotene, lutein, phytofluene, bixin and phytoene.
- 25. (Currently amended) The method according to any of claims 22 to 24 claim 22, wherein the at least one carotenoid is astaxanthin, zeaxanthin, bixin or phytoene.
- 26. (Currently amended) The method according to any of claims 22-25 claim 22, wherein sterilization and cell disruption are carried out using steam or microwave radiation.
- 27. (Currently amended) The method according to any of claims 22-26 claim 22, wherein the carotenoids are extracted from the biomass using dichloromethane or supercritical carbon dioxide or tetrahydrofuran.
- 28. (Original) The method according to claim 27, wherein the carotenoids dissolved in supercritical carbon dioxide are isolated directly or are taken up in dichloromethane.
- 29. (Currently amended) The method according to any of claims 22-28 claim 22, wherein the carotenoids are extracted from the biomass in a one-stage or, if appropriate, multistage process.
- 30. (Currently amended) The method according to any of claims 22-29 claim 22, wherein solvents are removed from the biomass in step IA1) IIA1) using steam distillation.

31. (Currently amended) The method according to any of claims 22-30 claim 22, wherein drying in step IIA3) is carried out using spray drying or contact drying.

- 32. (Currently amended) The method according to any of the preceding claims claim 22, wherein crystallization in step IIB1) is carried out by replacing the solvent gradually with a solvent in which carotenoids are not soluble.
- 33. (Currently amended) The method according to claim 32, wherein the solvent used is replaced with water or with a lower alcohol, in particular methanol.
- 34. (Currently amended) The method according to claim 13, wherein the genetically modified organism of the Blakeslea genus can be produced by transformation with a vector which has comprises a sequence selected from the group consisting of SEQ ID NO: SEQ ID NOs: 37 51 and SEQ ID NO: 62.
- 35. (Currently amended) A method for producing a foodstuff comprising organisms of the Blakeslea genus and at least one carotenoid, which method comprises, after cultivation of carotenoid-producing genetically modified organisms of the Blakeslea genus according to any of claims 1 to 21 claim 1, the following steps:
 - I) homogeneous suspension of the solids of the culture broth, and
 - IIA) for a biomass solid content of the culture broth of > 2%:
 - optional concentration of the culture broth to give a solid content of
 < 50%, and
 - 2) drying of the culture broth to produce the foodstuff,
 - IIB) for a solid content of < 2% of the culture broth,

or

or

- 1) concentration of the culture broth to give a solid content of > 2% and < 50%, and
- 2) drying of the suspension to produce the foodstuff,
- IIC) independently of the solid content of the culture broth,

1) removal of the biomass,

- 2) optional washing of the biomass with solvents in which carotenoids are not soluble, in particular water,
- 3) sterilization and cell disruption,
- 4) optional drying and homogeneous distribution,
- 5) partial extraction of the carotenoids from the biomass using a carotenoid-dissolving solvent,
 - 5a) removal of the carotenoid-containing biomass from the carotenoid-containing solvent,
 - 5b) removal of residual solvent from the biomass, and
 - 5c) drying of the biomass to produce the foodstuff,
- 6) crystallization of the carotenoids from the solvent used in 5a) and isolation of the carotenoid crystals, in particular by filtration.
- 36. (Original) The method according to claim 35, wherein the at least one carotenoid is selected from the group consisting of carotenes and xanthophylls.
- 37. (Currently amended) The method according to claim 35 or 36, wherein the at least one carotenoid is selected from the group consisting of astaxanthin, zeaxanthin, echinenone, β-cryptoxanthin, andonixanthin, adonirubin, canthaxanthin, 3-hydroxyechinenone, 3'-hydroxyechinenone, lycopene, β-carotene, lutein, bixin and phytoene.
- 38. (Currently amended) The method according to any of claims 35-37 claim 35, wherein the at least one carotenoid is astaxanthin, zeaxanthin, bixin or phytoene.
- 39. (Currently amended) The method according to any of claims 35-38 claim 35, wherein sterilization and cell disruption in step H3) IIC3) are carried out using steam or microwave radiation.
- 40. (Currently amended) The method according to any of claims 35-39 claim 35, wherein the carotenoids are extracted from the biomass in step IIC5) using dichloromethane or supercritical carbon dioxide.

41. (Original) The method according to claim 40, wherein the carotenoids dissolved in supercritical carbon dioxide are isolated directly or are taken up in dichloromethane.

- 42. (Currently amended) The method according to any of claims 35-41 claim 35, wherein the carotenoids are extracted from the biomass in a one-stage or, if appropriate, multistage process.
- 43. (Currently amended) The method according to any of claims 35-42 claim 35, wherein solvents are removed from the biomass in step IIC5b) using steam distillation.
- 44. (Currently amended) The method according to any of claims 35-43 claim 35, wherein drying in any of steps HA1) IIA2), IIB2) and IIC5c) is carried out using spray drying or contact drying.
- 45. (Currently amended) The method according to any of claims 35-44 claim 35, wherein crystallization in step IIC6) is carried out by replacing the solvent gradually with a solvent in which carotenoids are not soluble.
- 46. (Currently amended) The method according to claim 45, wherein the solvent used is replaced with water or with a lower alcohol, in particular methanol.
- 47. (Currently amended) The method according to any of claims 35-46 claim 35, wherein the genetically modified organism of the Blakeslea genus can be produced by transformation with a vector which has a sequence selected from the group consisting of SEQ ID NO: SEQ ID NOs: 37 51 and SEQ ID NO: 62.
- 48. (Currently amended) A foodstuff, in particular animal feedstuff, which can be produced by any of the methods of claims 1 to 47 the method of claim 1.
- 49. (Currently amended) A food supplement, in particular animal feed supplement, which can be produced by any of the methods of claims 1 to 47 the method of claim 1.

50. (Currently amended) A method according to any of claims 1-49, wherein for producing the foodstuff and animal feedstuff of claim 48 can be obtained from comprising a fermentation.

- 51. (Currently amended) The A method according to any of claims 1-49, wherein for producing the food supplement and animal feed supplement of claim 49 can be obtained from comprising a fermentation.
- 52. (Currently amended) The method according to any of claims 1-49 claim 35, wherein at least two products of the group consisting of foodstuff, food supplement, animal feedstuff and animal feed supplement can be obtained from a fermentation.
- 53. (Currently amended) The use of method according to claim 35, wherein the carotenoids obtainable by a method of claims 1 to 14 for producing are incorporated in the production of cosmetic, pharmaceutical, dermatological preparations, foodstuffs, food supplements, animal feedstuff or animal feed supplement.
- 54. (New) The method according to claim 8, wherein the hygromycin resistance gene (hph) is from *E. coli*.
- 55. (New) The method according to claim 33, wherein the solvent is replaced with methanol.
- 56. (New) The method according to claim 46, wherein the solvent is replaced with methanol.